

NEW METHODS IN CONSERVATIVE TREATMENT OF SPLEEN INJURY IN CLOSED ABDOMINAL TRAUMA

Kurbanov Nizom Azizovich., Achilov Mirzakarim Temirovich., Mavlonov Azimjon Dilshodovich

*Samarkand Medical University, Samarkand, Uzbekistan
Samarkand branch of RNCMP*

Annatation

The spleen has a number of important functions, the main of which are participation in hematopoiesis, the body's immune status, and an important role in the hemostasis system [2, 5]. In particular, after splenectomy, there is a change in the main indicators of the vascular-platelet hemostasis link: an increase in the number of platelets and their functional activity, including adhesive ability, the reaction of the release of platelet factors is disturbed, and the retraction index decreases [7]. Changes can also be traced in the coagulation link of the hemostasis system: there is a shortening of the activated partial thromboplastin time, an increase in the concentration of fibrinogen, and inhibition of fibrinolysis. All this leads to the development of thrombotic complications. Under our supervision there were 25 victims with damage to the spleen with a combined injury of the abdomen, who underwent conservative hemostatic therapy. They were combined with skull trauma in 8 (32.0%) cases, chest - in 10 (40.0%) cases, pelvis and limbs - in 7 (28.0%) cases. Of these, there were 19 men (76.0%), women - 6 (24.0%), the average age was 29.5 ± 1.5 years. The causes of the combined injury of the spleen were: a fall from a height - in 13 (52.0%) patients, a car injury - in 12 (48.0%) patients. Terms of hospitalization of patients with spleen injury from the moment of injury: up to 1 hour - 18 (72.0%), from 1 to 3 hours - 7 (28%) victims. car injury - in 12 (48.0%) patients. Terms of hospitalization of patients with spleen injury from the moment of injury: up to 1 hour - 18 (72.0%), from 1 to 3 hours - 7 (28%) victims. car injury - in 12 (48.0%) patients. Terms of hospitalization of patients with spleen injury from the moment of injury: up to 1 hour - 18 (72.0%), from 1 to 3 hours - 7 (28%) victims.

The main criterion for choosing conservative treatment of patients was the presence of stable hemodynamics. All patients at admission, in addition to the clinical examination, underwent ultrasound every 6-8 hours in dynamics, multislice computed tomography (MSCT) of the abdominal cavity, a plain radiograph of the abdominal cavity and other anatomical regions by localization, a general blood and urine test, and a coagulogram.

Relevance. The spleen has a number of important functions, the main of which are participation in hematopoiesis, the body's immune status, and an important role in the hemostasis system [2, 5]. In particular, after splenectomy, there is a change in

the main indicators of the vascular-platelet hemostasis link: an increase in the number of platelets and their functional activity, including adhesive ability, the reaction of the release of platelet factors is disturbed, and the retraction index decreases [7]. Changes can also be traced in the coagulation link of the hemostasis system: there is a shortening of the activated partial thromboplastin time, an increase in the concentration of fibrinogen, and inhibition of fibrinolysis. All this leads to the development of thrombotic complications [8].

The participation of the spleen in the immune status of the body has been proven. The spleen belongs to the peripheral lymphoid organs [3,4]. It concentrates suppressor, helper lymphocytes and part of effector cells, as well as the process of active antibody formation and production of humoral mediators of immunity [4]. The spleen contains approximately 35% T-lymphocytes and about 65% B-lymphocytes [3, 6]. Both stages of differentiation of antibody-forming cells from bone marrow precursors take place in it, while for T-lymphocytes, the antigen-independent stage of differentiation from bone marrow precursors occurs in the thymus, and the antigen-dependent stage occurs in the spleen [10]. The complex structure of the lymphatic follicles of the spleen, including thymus-dependent, thymus-independent and macrophage elements, creates favorable conditions in the body for cell cooperation in the immune response [1,6]. There is no doubt that the removal of the spleen leads to a change in the immune status of the body and the development of infectious complications, especially those caused by opportunistic microflora.

In addition, splenectomy leads to the development of severe purulent-septic complications, both in the immediate and late postoperative period. It is known that in the immediate postoperative period after splenectomy, the number of postoperative complications reaches 30% [12]. In this case, lethality is 16-30% [11].

In order to prevent the development of complications and preserve the main functions of the organ, various types of organ-preserving operations are currently proposed, if it is impossible to use them, autolien transplantation of fragments of the damaged spleen [9].

At present, with the development and improvement of laparoscopic techniques and imaging equipment, a new opportunity has appeared in the diagnosis and conservative treatment of spleen injuries. But the paucity of works in the periodical literature devoted to the study of non-surgical treatment of spleen injuries to preserve the organ urgently requires scientific research in this direction.

The purpose of the study. There was an improvement in the results of complex

treatment for spleen injuries with closed abdominal trauma.

Material and methods. Under our observation there were 50 patients with spleen injury due to combined abdominal trauma, who underwent conservative hemostatic therapy. They were combined with skull trauma in 8 (32.0%) cases, chest - in 10 (40.0%) cases, pelvis and limbs - in 7 (28.0%) cases. Of these, there were 19 men (76.0%), women - 6 (24.0%), the average age was 29.5 ± 1.5 years. The causes of the combined injury of the spleen were: a fall from a height - in 13 (52.0%) patients, a car injury - in 12 (48.0%) patients. Terms of hospitalization of patients with spleen injury from the moment of injury: up to 1 hour - 18 (72.0%), from 1 to 3 hours - 7 (28%) victims.

The main criterion for choosing conservative treatment of patients was the presence of stable hemodynamics. All patients at admission, in addition to the clinical examination, underwent ultrasound every 6-8 hours in dynamics, multislice computed tomography (MSCT) of the abdominal cavity, a plain radiograph of the abdominal cavity and other anatomical regions by localization, a general blood and urine test, and a coagulogram.

When looking for fluid during ultrasound, the lowest-lying areas of the abdomen were examined in all projections. The fluid was visualized as an anechoic zone. Small amounts of fluid were collected from women in the retrouterine space (in the space of Douglas), in men - in the hepatorenal recess (in the pocket of Morrison).

Subsequently, the tactics of management were determined depending on the severity of the patient's condition, the volume of hemoperitoneum, the intensity of blood loss (BCC), hemodynamic parameters (CVP, heart rate, diuresis).

Results. All patients with spleen injury upon admission to the hospital showed clear signs of intraperitoneal bleeding: abdominal pain, collapse, positive symptoms of Kerr, Eleker "Roly-Vstanki", Kulenkamph. In ultrasound examination, the volume of hemoperitoneum ranged from 200 to 350 ml in 15 patients, from 350 ml to 500 ml in 9, and more than 1200 ml in 1. The latter underwent blood removal by video laparoscopy with a favorable outcome.

All patients required transfusion of cryoprecipitate of the corresponding blood group, on average, 3.5 ± 0.3 doses, 16 patients - fresh frozen plasma, 350.0 ± 55.8 ml each patient and 9 patients - 250 ± 35.0 ml of donor erythrocyte mass. By the 7th day of treatment, ultrasound and MSCT revealed no free fluid in the abdominal cavity. As experience gained, we determined the criteria for conservative treatment: no signs of continued intraperitoneal bleeding; stable hemodynamic parameters (BP, CVP, heart rate) against the background of ongoing infusion and hemostatic therapy;

the amount of blood poured into the abdominal cavity up to 400-500 ml (<10% BCC); no signs of peritonitis and damage to other abdominal organs.

The methods of conservative adequate hemostasis used by us in spleen injuries were effective in 93.3% of cases. There were no complications and deaths due to the use of conservative therapy and video-laparoscopic technique.

Conclusions.

Non-surgical treatment of spleen injuries is a risky event and requires good equipment with visualization equipment in a medical institution, as well as the possibility of constant monitoring of the patient.

Therapeutic videolaparoscopy can become an alternative to laparotomy, which can reduce the incidence of postoperative complications.

Literature

1. Kurbonov N. A. et al. MANAGEMENT OF THE MIRIZZI SYNDROME AND THE SURGICAL IMPLICATIONS OF CHOLECYSTCHOLEDOCHAL FISTULA //INTERNATIONAL SCIENTIFIC REVIEW OF THE PROBLEMS OF NATURAL SCIENCES AND MEDICINE. – 2021. – С. 24-32.
2. Курбанов Н. А. и др. Differentiated surgical tactics for Mirizzi syndrome in patients with cholelithiasis //Шпитальна хірургія. Журнал імені ЛЯ Ковальчука. – №. 1. – С. 56-61.
3. Kurbonov N. A. et al. MANAGEMENT OF THE MIRIZZI SYNDROME AND THE SURGICAL IMPLICATIONS OF CHOLECYSTCHOLEDOCHAL FISTULA //INTERNATIONAL SCIENTIFIC REVIEW OF THE PROBLEMS OF NATURAL SCIENCES AND MEDICINE. – 2021. – С. 24-32.
4. Курбонов Н. А., Исматуллаев Ф. Х. ДИФФЕРЕНЦИРОВАННЫЙ ХИРУРГИЧЕСКИЙ ПОДХОД ПРИ СИНДРОМЕ МИРИЗЗИ //EUROPEAN RESEARCH: INNOVATION IN SCIENCE, EDUCATION AND TECHNOLOGY. – 2022. – С. 95-101.
5. Янгиев Б. А. и др. Тактические подходы в современном лечении травматических повреждений печени //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 1. – С. 204-210.
6. Нарзуллаев С. И. и др. Синдром внутрибрюшной гипертензии при сочетанных абдоминальных травмах //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 1. – С. 211-220.

7. Турсунов О. А. и др. В условиях экстренной медицинской помощи раствора кристаллических аминокислот для дополнительного парентерального питания в интенсивной терапии больных оперированных на желудочно-кишечном тракте //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 1. – С. 221-228.
8. Сабилов Б. и др. Лечебно-диагностическая видеолапароскопия в экстренной хирургии //Журнал проблемы биологии и медицины. – 2015. – №. 1 (82). – С. 70-72.
9. Курбонов Н. А., Карабаев Х. К., Нормаматов Б. П. ИЗМЕНЕНИЯ ЛИПИДНОГО ОБМЕНА ПРИ ОЖГОВОЙ БОЛЕЗНИ У ДЕТЕЙ //Детская хирургия. – 2020. – Т. 24. – №. S1. – С. 47-47.
10. Курбонов Н. А., Карабаев Н. А., Тагаев К. Р. ОЖГОВЫЙ ШОК У ДЕТЕЙ //Детская хирургия. – 2020. – Т. 24. – №. S1. – С. 48-48.
11. Курбонов Н. А., Давлатов С. С., Рахманов К. Э. OPTIMIZATION OF SURGICAL TREATMENT OF PATIENTS WITH MIRIZZI SYNDROME //ЖУРНАЛ ГЕПАТО-ГАСТРОЭНТЕРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ. – 2022. – Т. 3. – №. 1.
12. Кадыров Р. и др. Эндоскопические методы гемостаза при кровотечении из варикозно расширенных вен пищевода //Журнал проблемы биологии и медицины. – 2017. – №. 4 (97). – С. 44-47.
13. Кадыров Р. и др. Сочетанный эндоскопический гемостаз при язвенных кровотечениях //Журнал проблемы биологии и медицины. – 2018. – №. 1 (99). – С. 47-49.
14. Курбонов Н., Давлатов С., Амонов М. Modern methods of diagnosis and treatment of patients with Mirizzi syndrome //Журнал вестник врача. – 2021. – Т. 1. – №. 2 (99). – С. 158-161.
15. QURBONOV N. A. et al. Current trends in the management of Mirizzi syndrome: A review of literature //Annals of the Romanian Society for Cell Biology. – 2021. – С. 1927-1932.
16. Abduraxmanovich A. A. et al. POSSIBILITIES OF APPLYING MINIMALLY INVASIVE TECHNOLOGIES IN TREATMENT OF SEVERE ACUTE PANCREATITIS //Central Asian Journal of Medical and Natural Science. – 2021. – Т. 2. – №. 3. – С. 246-249.
17. Курбонов Н. А. и др. MODERN APPROACHES TO THE TREATMENT OF DEEP BURNING PATIENTS //УЗБЕКСКИЙ МЕДИЦИНСКИЙ ЖУРНАЛ. – 2022. – Т. 3. – №. 2.
18. Davlatov S. et al. Current State of The Problem Treatment of Mirizzi Syndrome (Literature Review) //International Journal of Pharmaceutical Research. – 2020. – Т. 12. – №. Suppl. ry 2. – С. 1931-1939.

19. Qurbonov N. A., Davlatov S. S., Amonov M. M. MODERN METHODS OF DIAGNOSIS AND TREATMENT OF PATIENTS WITH MIRIZZI SYNDROME.
20. Azizovich Q. N., Axmedovich Y. B., Salim D. MIRIZZI SYNDROME: HISTORY, CURRENT KNOWLEDGE AND PROPOSAL OF A SIMPLIFIED CLASSIFICATION //Surgery. – 1950. – Т. 132. – С. 300.
21. Курбонов Н. А., Исматуллаев Ф. Х. ДИФФЕРЕНЦИРОВАННЫЙ ХИРУРГИЧЕСКИЙ ПОДХОД ПРИ СИНДРОМЕ МИРИЗЗИ //EUROPEAN RESEARCH: INNOVATION IN SCIENCE, EDUCATION AND TECHNOLOGY. – 2022. – С. 95-101.
22. Ro'ziyev P. N. et al. XOLETSISTIT VA UNING ASORATLARI BILAN OG'RIGAN BEMORLARNI DIAGNOSTIKA VA DAVOLASHNING ZAMONAVIY USULLARI //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 4. – С. 26-32.
23. Qurbonov N. A. et al. SURUNKALI PANKREATITNI DIAGNOSTIKA QILISH VA DAVOLASHNING ZAMONAVIY USULLARI //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 4. – С. 33-41.
24. Qurbonov N. A. et al. O'TKIR PANKREATITNI DIAGNOSTIKA QILISH VA DAVOLASHNING ZAMONAVIY USULLARI //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 4. – С. 42-52.
25. Qurbonov N. A. et al. O'TKIR PANKREATITNI DIAGNOSTIKA QILISH VA DAVOLASHNING ZAMONAVIY USULLARINI OPTIMALLASHTIRISH //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 4. – С. 53-59.
26. Qurbonov N. A. et al. SURUNKALI VA O 'TKIR PANKREATITNI DIAGNOSTIKA VA DAVOLASHNING ZAMONAVIY VA YANGI USULLARI //Journal the Coryphaeus of Science. – 2023. – Т. 5. – №. 4. – С. 60-67.
27. Эргашева Ф., Икромов Ш., Курбонов Н. А. НОВЫЕ МЕТОДЫ ЛЕЧЕНИЯ ЖЕНЩИН В ПЕРИОД КЛИМАКТЕРИЧЕСКОГО СИНДРОМА //INNOVATIVE DEVELOPMENTS AND RESEARCH IN EDUCATION. –2023. –Т. 2. –№. 14. –С. 252-257.